

Smoking Attributable Death Statistics per 100,000 Population

Note: Information can be compared/ranked by state.

State	Smoking Attributable Deaths per 100,000 Population	Smoking Attributable Lung Cancer Deaths per 100,000 Population	Smoking Attributable Respiratory Disease Deaths per 100,000 Population
Alabama	317.5	101.8	81.6
Alaska	270.4	89.4	75.7
Arizona	247.4	73.9	77.0
Arkansas	323.7	110.0	80.5
California	235.0	68.9	70.6
Colorado	237.6	64.1	85.4
Connecticut	238.3	77.6	61.8
Delaware	280.9	96.9	67.7
District of Columbia	249.9	85.6	43.3
Florida	258.8	87.1	64.4
Georgia	299.4	95.8	79.7
Hawaii	167.6	54.0	32.3
Idaho	237.4	68.1	76.6
Illinois	263.1	87.6	63.6
Indiana	308.9	102.2	83.6
Iowa	248.0	80.6	69.6
Kansas	262.7	84.1	76.6
Kentucky	370.6	128.6	96.6
Louisiana	299.8	105.3	66.4
Maine	289.8	97.1	85.6
Maryland	261.9	88.8	64.2
Massachusetts	249.4	84.7	66.2
Michigan	281.9	89.3	71.0
Minnesota	215.1	72.4	59.0
Mississippi	333.6	109.0	80.1
Missouri	307.8	101.2	78.5
Montana	276.0	83.6	93.1
Nebraska	235.8	75.4	70.8
Nevada	343.7	100.1	107.3
New Hampshire	272.4	86.0	76.9
New Jersey	239.5	79.7	54.0

Smoking Attributable Death Statistics per 100,000 Population

Note: Information can be compared/ranked by state.

State	Smoking Attributable Deaths per 100,000 Population	Smoking Attributable Lung Cancer Deaths per 100,000 Population	Smoking Attributable Respiratory Disease Deaths per 100,000 Population
New Mexico	234.0	60.8	77.6
New York	246.1	74.6	57.8
North Carolina	298.4	96.5	78.2
North Dakota	225.6	70.1	60.7
Ohio	299.1	96.4	79.4
Oklahoma	332.1	101.2	89.9
Oregon	263.3	87.8	76.9
Pennsylvania	259.0	84.7	62.1
Rhode Island	266.8	88.9	65.9
South Carolina	293.4	96.3	73.6
South Dakota	239.2	74.9	67.6
Tennessee	325.0	108.6	85.7
Texas	273.1	85.2	72.9
Utah	138.3	34.7	48.9
Vermont	247.5	79.0	74.9
Virginia	267.0	89.6	70.3
Washington	261.0	85.7	75.1
West Virginia	344.3	110.8	96.0
Wisconsin	244.2	76.5	65.1
Wyoming	283.1	75.4	103.6

Health impact information is taken from the Smoking Attributable Mortality, Morbidity and Economic Costs (SAMMEC) software. Smoking attributable death rates reflect average annual estimates for the period 2000-2004, are calculated for persons aged 35 years and older and are age-adjusted to the 2000 U.S. population. They do not take into account deaths from burns or secondhand smoke. Respiratory diseases include pneumonia, influenza, bronchitis, emphysema and chronic airway obstruction.